

A Spectacular Radio Flare from XRF 050416a at 40 days and Implications for the Nature of X-ray Flashes

A. M. Soderberg¹, E. Nakar², S. B. Cenko³, P. B. Cameron¹, D. A. Frail⁴, S. R. Kulkarni¹, D. B. Fox⁵, E. Berger^{6,7,8}, A. Gal-Yam^{1,8}, D-S. Moon³, P. A. Price⁹, G. Anderson¹⁰, B. P. Schmidt¹⁰, M. Salvo¹⁰, J. Rich¹⁰, A. Rau¹, E. O. Ofek¹, R. A. Chevalier¹¹, M. Hamuy⁶, F. A. Harrison³, P. Kumar¹², A. MacFadyen¹³, P. J. McCarthy⁶, H. S. Park¹⁴, B. A. Peterson¹⁰, M. M. Phillips⁶, M. Rauch⁶, M. Roth⁶, S. Sheckman⁶

ABSTRACT

We present detailed optical, near-infrared, and radio observations of the X-ray flash 050416a obtained with Palomar and Siding Springs Observatories as well as the *Hubble Space Telescope* and Very Large Array, placing this event among the best-studied X-ray flashes to date. In addition, we present an optical spectrum from Keck LRIS from which we measure the redshift of the burst, $z = 0.6528$. At this redshift the isotropic-equivalent prompt energy release was about 10^{51} erg,

¹Division of Physics, Mathematics and Astronomy, 105-24, California Institute of Technology, Pasadena, CA 91125

²Theoretical Astrophysics, 130-33 California Institute of Technology, Pasadena, CA 91125

³Space Radiation Laboratory, MS 220-47, California Institute of Technology, Pasadena, CA 91125

⁴National Radio Astronomy Observatory, Socorro, NM 87801

⁵Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802

⁶Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101

⁷Princeton University Observatory, Peyton Hall, Ivy Lane, Princeton, NJ 08544

⁸Hubble Fellow

⁹Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

¹⁰Research School of Astronomy and Astrophysics, Australian National University, Mt Stromlo Observatory, via Cotter Rd, Weston Creek, ACT 2611, Australia

¹¹Department of Astronomy, University of Virginia, P.O. Box 3818, Charlottesville, VA 22903-0818

¹²Astronomy Department, University of Texas, Austin, TX 78731

¹³Institute for Advanced Study, Princeton, NJ 08540

¹⁴Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, CA 94550

and using a standard afterglow synchrotron model we find that the blastwave kinetic energy is a factor of 10 larger, $E_{K,\text{iso}} \approx 10^{52}$ erg. The lack of an observed jet break to $t \sim 20$ days indicates that the opening angle is $\theta_j \gtrsim 7^\circ$ and the total beaming-corrected relativistic energy is $\gtrsim 10^{50}$ erg. We further show that the burst produced a strong radio flare at $t \sim 40$ days accompanied by an observed flattening in the X-ray band which we attribute to an abrupt circumburst density jump or an episode of energy injection (either from a refreshed shock or off-axis ejecta). Late-time observations with the *Hubble Space Telescope* show evidence for an associated supernova with peak optical luminosity roughly comparable to that of SN 1998bw. Next, we show that the host galaxy of XRF 050416a is actively forming stars at a rate of at least $2 \text{ M}_\odot \text{ yr}^{-1}$ with a luminosity of $L_B \approx 0.5L^*$ and metallicity of $Z \sim 0.2 - 0.8 Z_\odot$. Finally, we discuss the nature of XRF 050416a in the context of short-hard gamma-ray bursts and under the framework of off-axis and dirty fireball models for X-ray flashes.

Subject headings: gamma-ray bursts: specific (XRF 050416a)

1. Introduction

Nearly a decade ago, X-ray flashes (XRFs) were observationally recognized as a subclass within the sample of gamma-ray bursts (GRBs) detected by the *BeppoSAX* Wide-Field Cameras (Heise *et al.* 2001). The events are distinguished by a prompt spectrum that peaks in the soft X-ray range ($E_p \lesssim 25$ keV), a factor of ~ 10 below the typical values observed for GRBs (Band *et al.* 1993). Since then, it has been shown that XRFs and GRBs share many observational properties, including prompt emission durations (Sakamoto *et al.* 2005), redshifts (Soderberg *et al.* 2004a), broadband afterglows (e.g., XRF 050406, Romano *et al.* 2006; Schady *et al.* 2006), and host galaxy properties (Bloom *et al.* 2003; Jakobsson *et al.* 2004; Rau, Salvato & Greiner 2005). Moreover, the recent discovery of Type Ic supernovae (SNe) in association with XRFs 020903 (Soderberg *et al.* 2005b; Bersier *et al.* 2006) and 060218 (Pian *et al.* 2006; Mirabal *et al.* 2006; Modjaz *et al.* 2006) indicate that XRFs, like GRBs, are produced in massive stellar explosions. Together, these clues strongly suggest that XRFs and GRBs share similar progenitors.

Driven by this progress, several theories have been proposed to explain the soft prompt emission observed for XRFs under the framework of a standard GRB model. One popular idea posits that XRFs are merely typical GRBs viewed away from the collimation axis (e.g., Yamazaki, Yonetoku & Nakamura 2003). In this scenario the prompt emission is primarily beamed away from our line-of-sight, resulting in lower fluence and E_p values for the observed

burst. An important implication of the off-axis model is that the early afterglow evolution should be characterized by a rising phase as the jet decelerates and spreads sideways into our line-of-sight (Granot *et al.* 2002; Waxman 2004; Soderberg *et al.* 2006e).

Another theory suggests that XRFs are intrinsically different from GRBs in their ability to couple energy to highly-relativistic material. In this scenario, XRFs are produced in explosions characterized by lower bulk Lorentz factors, $10 \lesssim \Gamma \lesssim 100$, than those inferred for typical GRBs, $\Gamma \gtrsim 100$ (Zhang, Woosley & Heger 2004). This may be the result of baryon loading of the high-velocity ejecta, a so-called “dirty fireball” (Dermer, Chiang & Böttcher 1999). Generally speaking, low Lorentz factor explosions may be identified through an analysis of their prompt emission since an optically thin spectrum at high energies implies a lower limit on the Lorentz factor (Lithwick & Sari 2001). In the case of X-ray flashes, however, there are generally insufficient high energy photons for this type of analysis. For these events, detailed modeling of the broadband afterglow may be used to place a lower limit on the Lorentz factor.

Here we present an extensive, multi-frequency data set for XRF 050416a at $z = 0.6528$ which extends to $t \sim 220$ days after the burst. By combining near-infrared, optical, ultraviolet, radio and X-ray data we present an in-depth analysis of the afterglow, energetics, supernova and host galaxy of XRF 050416a, placing it among the best-studied X-ray flashes to date. Moreover, thanks to our dedicated late-time monitoring campaign, we show that XRF 050416a produced a strong radio flare at $t \sim 40$ days accompanied by a brief plateau phase in the X-ray band. Finally, we discuss the nature of XRF 050416a in the context of off-axis and dirty fireball models for X-ray flashes.

2. Observations

XRF 050416a was discovered by the *Swift* Burst Alert Telescope (BAT) on 2005 April 16.4616 UT. The ratio of 15–25 keV and 25–50 keV channel fluences, $f_{15-25 \text{ keV}}/f_{25-50 \text{ keV}} \approx 1.1$, classifies the event as an X-ray flash (Sakamoto *et al.* 2006). This is consistent with the low peak photon energy, $E_p = 15.0_{-2.7}^{+2.3}$ keV (Sakamoto *et al.* 2006); a factor of ~ 10 lower than the typical values observed for long-duration GRBs (Band *et al.* 1993) and a factor of ~ 3 larger than the values inferred for XRFs 020903 and 060218 (Sakamoto *et al.* 2004; Campana *et al.* 2006).

As discussed by Sakamoto *et al.* (2006), the prompt emission light-curve is characterized by a relatively smooth, triangular peak which is only detected at energies below 50 keV. The burst duration is $T_{90} \approx 2.4$ sec (15-150 keV), placing it between the classes of

short- and long-duration GRBs while the hardness ratio shows a clear softening. In addition, Sakamoto *et al.* (2006) note two intriguing features of the data: (1) the rise time of the pulse is longer than the decay time, and (2) the cross-correlation lag function (an indication of the spectral softening) is $-0.066^{+0.014}_{-0.018}$ sec, apparently inconsistent with the overall softening trend observed for the light-curves. This lag function estimate is significantly different than the typical values inferred for long-duration GRBs and even more extreme than the zero spectral lags inferred for short-hard bursts (Norris & Bonnell 2006). We note, however, that the spectral lag for XRF 050416a was derived through a comparison of the two softest BAT bands (15-25 and 25-50 keV) and therefore prevents a clear comparison with the spectral lag estimates for other *Swift* GRBs for which 15-25 and 50-100 keV bands are typically used. Given that the temporal evolution is not strongly variable, this may indicate that the prompt emission was produced by another process (e.g. external shocks, Dermer, Chiang & Böttcher 1999). Finally, we note that this burst is inconsistent with the lag-luminosity correlation for long-duration GRBs which posits that low luminosity bursts such as XRF 050416a have long spectral lags (Norris, Marani & Bonnell 2000).

2.1. Early Optical Observations

Using the roboticized Palomar 60-inch telescope (P60) we initiated observations of the field of XRF 050416a at 2005 April 16.4634 UT (2.5 minutes after the burst). In our first 120 second image we discovered a new source within the *Swift*/BAT error circle at $\alpha = 12^{\text{h}}33^{\text{m}}54^{\text{s}}.58$, $\delta = +21^{\circ}03'26''.7$ (J2000) with an uncertainty of 0.5 arcsec in each coordinate based on an astrometric tie to the USNO-B catalog (Figure 1). We subsequently monitored the afterglow evolution with the Palomar 60-inch, 200-inch, and Siding Springs 2.3-meter telescopes in the R -, I -, z' - and K_s -band through $t \approx 7$ days.

Aperture photometry was performed on each of the images in the standard method using the **apphot** package within IRAF. Absolute calibration of R -, I -, and z' -band light-curves was derived using the field calibration of Henden (2005) and utilizing the transformation equations of Smith *et al.* (2002). The K_s -band light-curve was calibrated against 2MASS using 15 unconfused sources. The errors resulting from calibration uncertainty ($\lesssim 10\%$) were added in quadrature to the measurement errors. As shown in Figure 2 and Table 1, the afterglow was $I = 18.82 \pm 0.11$ mag at $t \approx 1.6$ min (mid-exposure).

We supplement these NIR/optical afterglow data with additional measurements from the GCNs (Li *et al.* 2005; Price *et al.* 2005; Qiu *et al.* 2005; Torii 2005; Yanagisawa, Toda & Kawai 2005) and those reported by Holland *et al.* (2006), obtained with the *Swift*/UVOT and 1.54-m Danish Telescope. The resulting dataset spans 1930Å (*Swift*/UVOT UVW2) to 22000Å

(K_s); however, we note that the majority of the UV observations are upper limits. Using this extended dataset we measure the temporal and spectral properties of the NIR/optical afterglow emission. We find the following NIR/optical power-law decay indices (α , where $F_\nu \propto t^\alpha$) between ~ 0.01 and 1 day: $\alpha_{K_s} = -0.7 \pm 0.3$, $\alpha_I = -0.7 \pm 0.3$, $\alpha_R = -0.5 \pm 0.3$, $\alpha_V = -1.0 \pm 0.3$, $\alpha_B = -0.5 \pm 0.3$, consistent with the values reported by Holland *et al.* (2006). These values imply a mean temporal index of $\bar{\alpha}_{\text{NIR/opt}} = -0.7 \pm 0.2$.

Finally, we analyze the spectral index (β with $F_\nu \propto \nu^\beta$) within the NIR/optical bands. As shown in Figure 3 there are two epochs at which the photometric spectrum is well-sampled: $t \approx 0.014$ and 0.8 days. We fit each of the observed spectra with a simple power-law and find $\beta_{\text{NIR/opt}} \approx -1.3$ ($\chi_r^2 \approx 0.5$) and -1.5 ($\chi_r^2 \approx 0.6$) for the first and second epochs, respectively. As will be discussed in §3.1, the observed steep spectrum is indicative of extinction within the host galaxy.

2.2. Late-time Observations with HST

Using the *Wide-Field Camera (WFC)* of the *Advanced Camera for Surveys (ACS)* onboard the *Hubble Space Telescope (HST)*, we imaged the field of XRF 050416a four times, spanning 37 to 219 days after the burst. Each epoch consisted of two or four orbits during which we imaged the field in filters F775W and/or F850LP, corresponding to SDSS i' - and z' - bands, respectively.

The *HST* data were processed using the `multidrizzle` routine (Fruchter & Hook 2002) within the `stsdas` package of IRAF. Images were drizzled using `pixfrac=0.8` and `pixscale=1.0` resulting in a final pixel scale of 0.05 arcsec/pixel. Drizzled images were then registered to the final epoch using the `xregister` package within IRAF. We astrometrically tied the *HST* and P60 images using 12 unconfused sources in common resulting in a final systematic uncertainty of 0.70 arcsec (2σ).

To search for source variability and remove host galaxy contamination, we used the *ISIS* subtraction routine by Alard (2000) which accounts for temporal variations in the stellar PSF. Adopting the final epoch observations as templates we produced residual images. These residual images were examined for positive sources positionally coincident with the P60 afterglow.

Photometry was performed on the residual sources within a 0.5 arcsec aperture. We converted the photometric measurements to infinite aperture and calculated the corresponding AB magnitudes within the native *HST* filters using the aperture corrections and zero-points provided by Sirianni *et al.* (2005). Here we made the reasonable assumption that the tran-

sient flux is negligible in the template images. For comparison with ground-based data, we also converted the F775W measurements to Johnson *I*-band (Vega) magnitudes using the transformation coefficients derived by Sirianni *et al.* (2005) and assuming the F_ν source spectrum implied by the first epoch *HST* data.

As shown in Table 2 and Figure 4, the transient is clearly detected in the first epoch *HST* residual images. An astrometric tie between the first and final epochs shows that the residual is offset 0.02 ± 0.02 arcsec from the center of the host galaxy. Our residual images show the source to be $F775W = 24.35 \pm 0.02$ and $F850LP = 23.83 \pm 0.03$ mag in the AB system ($I = 23.82 \pm 0.02$ mag in the Vega system) at $t \approx 37$ days. As shown in Figure 2, these values are a factor of ~ 5 above an extrapolation of the early afterglow decay. The observed spectral index between the F775W and F850LP filters is $\beta_{HST} \approx -2.8 \pm 0.3$, significantly steeper than the afterglow spectrum observed at early time (§2.1 and Figure 3) as well as the typical values measured for NIR/optical afterglows ($\beta_{NIR/opt} \approx -0.6$; Panaitescu & Kumar 2002; Yost *et al.* 2003). As will be discussed in §4.4, the timescale and spectral signature of the observed flux excess are suggestive of a thermal supernova component.

2.3. Spectroscopic Observations

We observed the host galaxy of XRF 050416a with the Low Resolution Imaging Spectrograph on Keck I on 2005 June 6.3 UT ($t \sim 50$ days). We placed a 1.0 arcsec longslit across the host galaxy at a position angle of $PA=87^\circ$. Data were reduced in standard manner using the `onedspec` and `twodspec` packages within IRAF. Flux calibration was performed using the spectrophotometric standard star BD+284211.

As shown in Figure 5 and Table 3, we detect several strong emission lines in the spectrum including $H\beta$, $H\gamma$, $[O\ II]\lambda 3727$, and $[O\ III]\lambda\lambda 4959, 5006$ at a redshift of $z = 0.6528 \pm 0.0002$. Adopting the standard cosmological parameters ($H_0 = 71\text{ km s}^{-1}\text{ Mpc}^{-1}$, $\Omega_M = 0.27$, $\Omega_\Lambda = 0.73$), the isotropic gamma-ray energy release is $E_{\gamma,iso} \approx 1.2 \pm 0.2 \times 10^{51}$ erg (1 keV – 10 MeV; Sakamoto *et al.* 2006). Compared with typical long-duration bursts, the prompt energy release of XRF 050416a is a factor of ~ 100 lower (Frail *et al.* 2001; Bloom, Frail & Kulkarni 2003; Amati 2006 and references therein).

2.4. Radio Observations

We began observing XRF 050416a with the Very Large Array¹ (VLA) on 2005 April 16.49 UT ($t \approx 37$ min). No radio source was detected coincident with the optical position to a limit of $F_\nu < 122 \mu\text{Jy}$ at $\nu = 8.46$ GHz. However, further observations at $t \approx 5.6$ days revealed a new radio source with $F_\nu \approx 101 \pm 34 \mu\text{Jy}$ coincident with the optical and X-ray afterglow positions at $\alpha = 12^{\text{h}}33^{\text{m}}54.594^{\text{s}} \pm 0.002$, $\delta = 21^\circ03'26.27'' \pm 0.04$ (J2000) which we identify as the radio afterglow.

We continued to monitor the radio afterglow at 1.43, 4.86 and 8.46 GHz through $t \approx 140$ days (Table 4). All observations were taken in standard continuum observing mode with a bandwidth of 2×50 MHz. We used 3C286 (J1331+305) for flux calibration, while phase referencing was performed against calibrators J1221+282 and J1224+213. The data were reduced using standard packages within the Astronomical Image Processing System (AIPS).

As shown in Figure 6, the evolution of the radio afterglow is dissimilar from those of typical GRBs and inconsistent with a standard blastwave model. Between $t \sim 20$ and 40 days an abrupt rebrightening (factor of ~ 3) is observed at all radio bands with a temporal index steeper than $\alpha_{\text{rad}} \approx 0.9$. Following this radio flare the emission decays rapidly with an index of $\alpha_{\text{rad}} \lesssim -1.5$. The peak radio luminosity at 8.46 GHz, $L_{\text{rad}} \approx 1.3 \times 10^{31} \text{ erg s}^{-1} \text{ Hz}^{-1}$, is typical for GRBs (Frail *et al.* 2003), a factor of ~ 10 and 10^3 larger than those of XRFs 020903 and 060218, respectively (Soderberg *et al.* 2004a, 2006c), and between 10^2 and 10^6 times higher than the peak radio luminosities observed for optically-selected Type Ibc supernovae (Soderberg 2006). By $t \approx 105$ days the radio afterglow is no longer detected at any frequency.

2.5. X-ray Observations

The afterglow of XRF 050416a was observed with the *Swift* X-ray Telescope (XRT) beginning 1.2 minutes after the burst and continuing through $t \sim 69$ days, placing it among the best studied X-ray afterglows to date (Nousek *et al.* 2006; O’Brien *et al.* 2006). From their analysis of the XRT data, Mangano *et al.* (2006) report a spectral index of $\beta_X = -1.04^{+0.11}_{-0.05}$ and evidence for significant absorption in the host galaxy, $N_H = 6.8^{+1.0}_{-1.2} \times 10^{21} \text{ cm}^{-2}$, corresponding to $E(B - V)_{\text{rest}} = 1.2 \pm 0.2$ assuming the conversion of Predehl & Schmitt (1995)

¹The Very Large Array and Very Long Baseline Array are operated by the National Radio Astronomy Observatory, a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

and a Milky Way extinction curve (Pei 1992). As shown in Figure 7, at $t = 10$ hours the X-ray luminosity was $L_{X,\text{iso}}(t = 10 \text{ hrs}) \approx 2.3 \times 10^{45} \text{ erg s}^{-1}$, placing it at the lower edge of the observed distribution for GRBs (Berger, Kulkarni & Frail 2003; Freedman & Waxman 2001).

As discussed by Mangano *et al.* (2006) and shown in Figure 7, the early evolution of the X-ray afterglow can be characterized by three phases: (1) an initial steep decay (2) a flattening between $t \approx 7$ and 20 minutes during which the flux is roughly constant, and (3) a resumed decay through $t \sim 20$ days. These three phases have been shown to be ubiquitous among *Swift* X-ray afterglows (Nousek *et al.* 2006). However, at $t \sim 20$ days there is a second flattening which continues through $t \sim 40$ days. By 70 days the X-ray afterglow is no longer detected, implying a significant steepening to $\alpha_X \lesssim -1.8$ between the last two observations. We note that the timescale for the X-ray flattening and subsequent steep decay is coincident with the observed radio flare.

3. Properties of the Early Afterglow

Using the detailed multi-frequency observations of the XRF 050416a afterglow we can constrain the physical properties of the ejecta and the circumburst density. We adopt a standard relativistic blastwave model in which the afterglow emission is produced through the dynamical interaction of the ejecta with the surrounding medium (the forward shock, FS) with an additional component from shock heating of the ejecta (the reverse shock, RS). In this scenario, the total energy density is partitioned between relativistic electrons, ϵ_e , and magnetic fields, ϵ_B , while the thermal energy of the shocked protons accounts for the fraction remaining (see Piran 1999 for a review). The shocked electrons are accelerated into a power-law distribution, $N(\gamma) \propto \gamma^{-p}$ above a minimum Lorentz factor, γ_m . The emission resulting from the forward and reverse shock components is described by a synchrotron spectrum characterized by three break frequencies — the self-absorption frequency, ν_a , the characteristic frequency, ν_m , and the cooling frequency, ν_c — and a flux normalization, F_{ν_m} (Sari, Piran & Narayan 1998). In modeling the afterglow spectral and temporal evolution, we adopt the formalism of Granot & Sari (2002) for a relativistic forward shock expanding into a constant density circumburst medium.

3.1. Preliminary Constraints

In fitting the forward shock model to the afterglow data of XRF 050416a we use only observations between 0.014 and 20 days when the afterglow follows a simple power-law evolution. To constrain the spectrum of the forward shock, we first investigate the afterglow evolution in the optical and X-ray bands. As shown in Figure 7, the X-ray data between 0.014 and 20 days are reasonably fit with $\alpha_X \approx -1.1$ ($\chi_r^2 \approx 0.70$). Mangano *et al.* (2006) report that the X-ray spectral index on this same timescale is $\beta_X \approx -1.04$, leading to $\alpha - 3\beta/2 \approx 0.5$. A comparison to the standard closure relations, $\alpha - 3\beta/2 = 0$ ($\nu_m < \nu < \nu_c$) and $\alpha - 3\beta/2 = 1/2$ ($\nu > \nu_c$) indicates that $\nu_X > \nu_c$. This conclusion is supported by the near-IR to X-ray spectral slope, $\beta_{K,X} = -0.47 \pm 0.06$ at $t \approx 0.6$ days, which is flatter than β_X as expected if $\nu_{\text{NIR/opt}} < \nu_c < \nu_X$. Therefore, the X-ray observations suggest that $p = -2\beta \approx 2.1$.

Next we consider the spectral index within the NIR/optical bands. As discussed in §2.1 and shown in Figure 3, the observed NIR/optical spectral index on this timescale is $\beta_{\text{NIR/opt}} \approx -1.3$ to -1.5 . These values are significantly steeper than $\beta_{K,X}$ and imply that the optical flux is suppressed by host galaxy extinction. Making the reasonable assumption that $\nu_m \lesssim \nu_{\text{NIR/opt}}$ on the timescale of our afterglow observations, and adopting $p \approx 2.1$ as indicated by the X-ray observations, we estimate $\beta_{\text{NIR/opt}} = -(p - 1)/2 \approx -0.55$ for the intrinsic spectral index of the NIR/optical afterglow. Adopting this value for $\beta_{\text{NIR/opt}}$, we find that both NIR/optical spectra are reasonably fit with a Galactic extinction of $E(B - V) = 0.03$ (Schlegel, Finkbeiner & Davis 1998) and a host galaxy component of $E(B - V)_{\text{rest}} \approx 0.28$ (Figure 3). Here we have assumed a Milky Way extinction model for the host (Pei 1992) but note that a comparable fit may be obtained for an SMC extinction curve. We further note that this optically-derived extinction estimate is lower than that inferred from the X-ray spectrum (Mangano *et al.* 2006), consistent with the trend observed for long-duration GRBs Galama & Wijers (2001).

With this extinction correction, the near-IR to X-ray spectral index becomes $\beta_{K,X} \approx -0.5 \pm 0.1$, consistent with our estimate for the intrinsic spectral index within the NIR/optical band. Moreover, the extinction-corrected NIR/optical spectral index and observed average temporal index of $\bar{\alpha}_{\text{NIR/opt}} = -0.7 \pm 0.2$ are consistent with the standard closure relation: $\alpha - 3\beta/2 = 0 \approx 0.1$. We also note that this supports our assumption of a constant density medium since in a wind environment the expected temporal index is steeper than $\alpha = -1.25$, and thus inconsistent with the observed values. Using all the available optical and X-ray observations we estimate that $\nu_c \approx 1 \times 10^{17}$ Hz at $t = 1$ day.

Next we compare the near-IR and radio afterglow data to constrain ν_m and the peak spectral flux, F_{ν_m} . Assuming that ν_m passed through the NIR/optical bands near the time of

our first K_s -band observations implies that the peak spectral flux is roughly comparable to the extinction-corrected K_s flux: $F_{K_s} \approx 230 \mu\text{Jy}$ at $t \approx 11$ minutes. Here we focus on the K_s -band data since they are the least sensitive to host galaxy extinction, which we estimate to be $A_K \approx 0.24$ mag (a 20% increase in flux) for $E(B-V)_{\text{rest}} \approx 0.28$. Scaling these constraints to $t = 1$ day ($\nu_m \propto t^{-1.5}$ and $F_{\nu_m} \propto t^0$) and accounting for the smooth shape of the spectral peak, we find $\nu_m \approx 4.0 \times 10^{11}$ Hz and $F_{\nu_m} \approx 350 \mu\text{Jy}$. Here and throughout, F_{ν_m} is the asymptotic extrapolation of the smooth spectrum peak and is therefore slightly higher than the intrinsic peak flux. We note that since the NIR/optical data require $F_{\nu_m} \propto \nu^{-(p-1)/2}$, lower values of ν_m imply increasingly higher values of F_{ν_m} at the time of the first K_s -band observations.

Finally, we test that these constraints are consistent with the radio observations. Given the evolution of ν_m , these constraints predict that the spectral peak should pass through the radio band at $t \approx 13$ days with an extrapolated peak flux density of $F_{\nu_m} \approx 350 \mu\text{Jy}$, roughly consistent with the 4.86 GHz observations on this timescale. We emphasize that the early steady decay of the NIR/optical data require that ν_m passes through the radio no later than 13 days. Finally, we note that the radio spectrum is optically thin throughout the timescale of VLA monitoring (see Figure 6) and thus we observationally constrain ν_a to be below 1.43 GHz.

3.2. Forward Shock Broadband Model

Adopting these constraints we apply a broadband afterglow model fit to the multi-frequency data in order to determine the physical parameters of the burst. The four spectral parameters (F_{ν_m} , ν_a , ν_m and ν_c) are fully determined by four physical parameters: the isotropic ejecta energy, $E_{K,\text{iso}}$, the energy density partition fractions, ϵ_e and ϵ_B , and the circumburst density, n . Therefore by constraining the four spectral parameters through broad-band observations, we are able to determine a unique solution for the four physical parameters (see Sari, Piran & Narayan 1998 and Piran 1999 for reviews). Although the radio observations provide only an upper limit on ν_a , we are able to define a range of reasonable values by requiring that $\epsilon_e, \epsilon_B \leq 1/3$ which accounts for an equal or greater contribution from shocked protons. This requirement excludes unphysical solutions in which the sum of the contributions from shocked electrons, protons and magnetic fields exceed the total energy density. Combined with the observed constraints for F_{ν_m} , ν_m and ν_c we find the following ranges for the physical parameters:

$$E_{K,\text{iso}} \approx (8.2 - 14) \times 10^{51} \text{ erg} \quad (1a)$$

$$n \approx (0.33 - 4.2) \times 10^{-3} \text{ cm}^{-3} \quad (1b)$$

$$\epsilon_e \approx (0.20 - 1/3) \quad (1c)$$

$$\epsilon_B \approx (0.072 - 1/3). \quad (1d)$$

As shown in Figures 2, 6, and 7, this model provides an adequate fit to the broadband data between $t \sim 0.01$ and 20 days.

3.3. Collimation of the Ejecta and Viewing Angle

The lack of an observed jet break in the X-rays to $t \sim 20$ days, together with the inferred physical parameters constrain the opening angle of the jet (e.g. Sari, Piran & Halpern 1999) to $\theta_j \approx 3.1 t_j^{3/8} E_{K,\text{iso},52}^{-1/8} n_{-3}^{1/8} (1+z)^{-3/8} \gtrsim 6.9$ degrees. Here, t_j is the jet break time in days and we have adopted the notation $10^x Q_x = Q$. This limit is slightly larger than the median of the jet opening angles inferred for long-duration GRBs, $\theta_j \sim 5^\circ$ (Bloom, Frail & Kulkarni 2003; Ghirlanda, Ghisellini & Lazzati 2004; Soderberg *et al.* 2006a and references therein). This indicates that the beaming-corrected ejecta energy release is $E_K \equiv E_{K,\text{iso}}(1 - \cos\theta) \approx (9.8 \times 10^{49} - 1.4 \times 10^{52})$ erg where the range includes the uncertainty in $E_{K,\text{iso}}$ and the lower limit on θ_j . Moreover, we expect the blastwave to become non-relativistic on a timescale $t_{\text{NR}} \approx 2.0 E_{51}^{1/3} n_{-3}^{-1/3} \sim (0.6 - 6.8)$ yrs (Livio & Waxman 2000). On a similar timescale, the ejecta are predicted to approach spherical symmetry after which the blastwave evolution is well described by the Sedov-von Neumann-Taylor (SNT) solution (Zel'dovich & Raizer 2002; Frail, Waxman & Kulkarni 2000); in this regime the afterglow emission decays with $\alpha = -9/10$ (-1) for frequencies below (above) the cooling frequency.

Next, the early steady decay of the X-ray and NIR/optical afterglow indicates that the jet collimation axis is directed roughly along our line-of-sight. In comparison, GRBs viewed significantly off-axis ($\theta_{\text{obs}} > 2\theta_j$) are predicted to show a rising afterglow light-curve as the jet spreads sideways and intersects our viewing angle (Granot *et al.* 2002; Waxman 2004; Soderberg *et al.* 2006e). Here, θ_{obs} is the angle between our line-of-sight and the jet collimation axis. We conclude that the ejecta are viewed roughly on-axis and therefore the inferred beaming-corrected energies are not affected significantly (if at all) by viewing angle effects.

4. Properties of the Late-time Afterglow

Next we address the nature of the late-time broadband afterglow evolution with special attention to the strong radio flare observed at ~ 40 days. Radio flares have been noted for several other GRBs, though only at early times (e.g. GRB 990123 at $t \lesssim 1$ day;

Kulkarni *et al.* 1999). Based on their observed timescale and evolution, radio flares are typically attributed to emission from the reverse shock (Sari & Piran 1999). Here we present detailed radio observations for XRF 050416a which show for the first time a strong radio flare at late time. Possible causes for a late-time radio rebrightening include the emission from a decelerating jet initially directed away from our line-of-sight (Waxman 2004; Li & Song 2004), circumburst density variations (Wijers 2001; Ramirez-Ruiz *et al.* 2001), and energy injection from a slow shell catching up to and refreshing the afterglow shock (Rees & Meszaros 1998). We discuss each of these possibilities below.

4.1. Off-axis Jet Emission

It has been shown that the observational signature of a relativistic jet viewed significantly away from our line-of-sight is a rapid achromatic rise in the early afterglow light-curves (Paczynski 2001; Granot *et al.* 2002; Waxman 2004). In this scenario, the observed peak of the afterglow emission occurs as the spreading jet crosses our viewing angle. The timescale for this peak is ~ 100 days for a GRB jet with typical parameters ($E_K = 10^{51}$ erg, $n = 1 \text{ cm}^{-3}$, $\epsilon_e = \epsilon_B = 0.1$, $\theta_j = 5^\circ$) viewed from an angle $\theta_{\text{obs}} = 30^\circ$ (Soderberg *et al.* 2006e). The subsequent afterglow evolution is the same as that seen by an on-axis observer, decaying steeply with $\alpha = -p \approx [-2 \text{ to } -3]$ for frequencies above ν_m (Sari, Piran & Halpern 1999). The observed timescale and evolution of the XRF 050416a radio flare are therefore roughly consistent with the predictions of an off-axis relativistic jet.

However, as noted by Mangano *et al.* (2006) and discussed in §3.3, the early and steady decay of the XRF 050416a broadband afterglow implies the presence of relativistic ejecta directed along our line-of-sight. These ejecta are also responsible for the production of the observed prompt emission. Attributing the strong late-time radio flare to an energetic off-axis relativistic jet therefore implies that multiple relativistic ejecta components were produced in the explosion. Moreover, the steep rise and peak flux of the radio flare imply sharp edges for the off-axis jet and a kinetic energy 2 to 3 times larger than that of the on-axis ejecta. While this scenario cannot be ruled out, we consider it unlikely given the complicated ejecta geometry required.

4.1.1. A Receding Jet

Li & Song (2004) describe a related scenario in which a strong late-time radio flare is observed from a receding jet initially directed anti-parallel to our line-of-sight. In this

case, early afterglow emission is expected from the approaching jet while emission from the receding jet is expected on a timescale $5 \times t_{\text{NR}}$ due to the light travel time delay. This scenario is appealing in that it may explain both the early and late-time afterglow emission observed for XRF 050416a within the standard framework of engine-driven (accretion-fed compact source; Piran 1999) double-sided jets. However, as discussed in §3.3, broadband modeling of the afterglow emission predicts the non-relativistic transition to occur no earlier than 0.6 yrs – too late to explain the radio flare at 40 days. Moreover, the peak emission from a receding jet should be comparable to the radio flux at t_{NR} and decay with a temporal index given by the Sedov solution, $\alpha_{\text{rad}} = -9/10$ (Li & Song 2004); both of these predictions are inconsistent with the observations. We conclude that afterglow emission from a receding jet is unlikely to produce the observed radio flare.

4.2. Circumburst Density Variations

It has been argued that abrupt variations in the circumburst medium can produce a strong rebrightening in the radio afterglows of GRBs. Specifically, the dynamical interaction of the forward shock with a wind-termination shock at $r \sim 1$ pc is predicted to cause a rebrightening of the radio afterglow on a timescale of a few years (Wijers 2001; Ramirez-Ruiz *et al.* 2001, 2005). Here it is assumed that the blastwave is expanding non-relativistically when it encounters the density jump, overall consistent with the observed timescales for non-relativistic transitions (Frail, Waxman & Kulkarni 2000; Berger, Kulkarni & Frail 2004; Frail *et al.* 2005). For comparison, the interaction of a relativistic blastwave with an abrupt (step function) density jump is not expected to cause strong variations in the afterglow light-curves, where here it has been assumed that the post-jump expansion is also relativistic (Nakar & Granot 2006).

In the case of XRF 050416a, the strong radio flare and X-ray plateau phase occur on a timescale when the blastwave is still relativistic, and therefore an abrupt circumburst density jump appears an unlikely explanation. This is supported by the fact that during the relativistic regime, the flux at frequencies above ν_c should be unaffected by circumburst density variations (Granot & Sari 2002). We speculate, however, that a very large density jump may be able to decelerate the blastwave to non-relativistic speeds on a very short timescale and may therefore be able to explain the unusual late-time afterglow evolution. The increase in density would cause a shift in ν_a which may explain the peculiar evolution of the spectral index in the radio band (Figure 6). Based on our afterglow modeling (§3.2), we estimate that the circumburst radius of the forward shock at the onset of the radio flare was roughly ~ 1 pc, roughly consistent with the radius of a wind-termination shock

(Garcia-Segura, Langer & Mac Low 1996; Chevalier, Li & Fransson 2004). In this scenario, we expect a post-jump self-similar evolution consistent with the Sedov-Taylor solution. Since the late-time radio and X-ray data are not sufficiently sensitive to trace the post-flare evolution, we cannot rule out this possibility.

Finally, we investigated a scenario where the radio/X-ray flare is produced by the dynamical interaction of the quasi-spherical, non-relativistic SN ejecta with a CSM density enhancement. In fact, density jumps have been invoked to explain radio modulations observed for local (non-relativistic) SNe Ibc (e.g. SN 2003bg; Soderberg *et al.* 2006b). Adopting a simple minimum energy calculation (Kulkarni *et al.* 1998) and requiring that the shock energy is equally partitioned between magnetic fields and relativistic electrons, we find that attributing the strong radio emission to the quasi-spherical SN component requires that the SN ejecta is relativistic ($\Gamma \sim 10$) at the time of the radio/X-ray flare. However, as discussed above, strong flux variations are not expected while the blastwave is relativistic. Combined with the fact that the flare is at least a factor of 10^2 more radio luminous than any other SN Ibc ever observed (including GRB-SN 1998bw, Kulkarni *et al.* 1998), we conclude that the radio/X-ray flare can not be attributed to the quasi-spherical SN ejecta.

4.3. Energy Injection

An episode of energy injection may also cause a rebrightening of the afterglow flux. Energy injection may arise from long-lived central engine activity or under the framework of a “refreshed” shock where a slow moving shell ejected during the initial burst eventually catches up with the afterglow shock (Rees & Meszaros 1998; Granot, Nakar & Piran 2003). The observed ratio of the settling time to the epoch of injection, $\delta t/t$, can distinguish between these two scenarios. For $\delta t/t < 1$ the injection is produced by engine activity, while $\delta t/t \gtrsim 1$ indicates a refreshed shock. In the case of XRF 050416a, the X-ray and radio data after 20 days imply $\delta t/t > 1$, suggesting that the ejecta were refreshed on this timescale.

As shown in Figure 6, an extrapolation of the early radio evolution lies a factor of ~ 3 below the observed flux at the onset of the radio flare. An energy increase affects the spectral parameters according to the following scalings: $\nu_a \propto E_{K,\text{iso}}^{1/5}$, $\nu_m \propto E_{K,\text{iso}}^{1/2}$, $F_{\nu_m} \propto E_{K,\text{iso}}$, and $\nu_c \propto E_{K,\text{iso}}^{-1/2}$ (Granot & Sari 2002). Given that ν_m is within the radio band on this timescale we have $F_{\nu,\text{rad}} \propto E_{K,\text{iso}}$ and thus the radio flare corresponds to an energy injection of a comparable factor, ~ 3 . For the X-ray band, $F_{\nu,X} \propto E_{K,\text{iso}}^{(p+2)/4}$, thus for $p \approx 2.1$ an energy injection of a factor of ~ 3 corresponds to a comparable increase in the X-ray flux. The *Swift*/XRT observations suggest a flattening on this timescale. Here we adopt the conservative assumption that the shock microphysics (ϵ_e , ϵ_B and p) do not evolve during the

energy injection.

One important implication of the energy injection model is that the post-injection asymptotic temporal decay should be the same as that before the injection. This prediction is consistent with the energy injection episodes invoked for GRBs 021004 and 030329 (Björnsson, Gudmundsson & Jóhannesson 2004; Granot, Nakar & Piran 2003). However, in the case of XRF 050416a, the late-time radio and X-ray data suggest a steep post-injection decay, $\alpha \gtrsim -2$, significantly steeper than that observed pre-injection. Moreover, the afterglow should asymptotically approach a flux normalization larger by a factor of ~ 3 in both the radio and X-ray bands. As shown in Figures 6 and 7, this appears inconsistent with the observations which suggest that the late-time afterglow is comparable (or fainter) than an extrapolation of the early afterglow model. We note, however, that the faintness of the late-time afterglow and the steep post-injection decay may be explained if a jet break occurred on roughly the same timescale as the energy injection.

In comparison with other late-time afterglow studies (Frail, Waxman & Kulkarni 2000; Tiengo *et al.* 2003; Berger, Kulkarni & Frail 2004; Kouveliotou *et al.* 2004; Frail *et al.* 2005), the radio flare and X-ray flattening observed for XRF 050416a are clearly atypical for long-duration GRBs. We therefore attribute the observed evolution to an unusual scenario involving either a large circumburst density jump or a late-time injection of energy (from a slow shell or off-axis ejecta).

4.4. An Associated Supernova

The *HST* measurements at late-time can be used to constrain any possible contribution from an associated supernova. Based on previous studies of GRB-SNe, the thermal emission from an associated supernova is predicted to reach maximum light on a timescale of $20 \times (1+z)$ days with a peak magnitude of $M_{V,\text{rest}} \approx -20$ mag or fainter (Zeh, Klose & Hartmann 2004; Soderberg *et al.* 2006d and references therein). Observationally, the emergence of a SN component produces a steepening of the optical spectrum as the supernova nears maximum light and dominates the afterglow emission.

An extrapolation of the broadband afterglow model to 40 days shows that the *HST* data are brighter by a factor of ~ 5 . For comparison, the radio flare and X-ray plateau on this same timescale represent flux density enhancements by factors of roughly ~ 3 and 2, respectively.

As discussed in §2.2 we measure the spectrum of the optical transient within the *ACS* bands and find $\beta_{HST} \approx -2.8 \pm 0.3$ at 37 days. After correction for extinction (see §2.1), the

implied spectral index becomes $\beta_{HST} \approx -1.9$. For comparison, the spectral index between the radio and X-ray bands on this timescale (coincident with the radio flare and X-ray flattening) is $\beta_{\text{radX}} \approx -0.56 \pm 0.04$ and the indices within the bands are $\beta_{\text{rad}} \approx -0.6 \pm 0.2$ (Figure 6) and $\beta_X \approx -1.0$ (Mangano *et al.* 2006), respectively. Given that the *HST* derived spectral index, even after correction for extinction, is (1) significantly steeper than the NIR/optical index observed at early time, $\beta_{\text{NIR/opt}} \approx -1.4$, (2) significantly steeper than β_{radX} measured at a comparable epoch, and (3) inconsistent with the range of synchrotron spectral indices predicted for a relativistic blastwave ($\beta = [-1.5, 2.5]$, Sari, Piran & Narayan 1998), we conclude that the optical flux at $t \sim 40$ days is dominated by another emission process, likely an associated SN.

Next, to determine if the late-time optical data are consistent with the temporal evolution of a typical GRB/XRF-associated supernova, we compare the *HST* flux values with synthesized SN light-curves. We compiled *UBVR_IJHK* observations of SN 1998bw from the literature (Galama *et al.* 1998; McKenzie & Schaefer 1999; Patat *et al.* 2001) and smoothed the extinction-corrected (Galactic component of $E(B-V)=0.059$, Schlegel, Finkbeiner & Davis 1998) light-curves. Here we assume that the SN 1998bw host-galaxy extinction is negligible, consistent with the findings of Patat *et al.* (2001) based on a spectroscopic analysis of the host galaxy. We then produced k -corrected NIR/optical light-curves of SN 1998bw at the redshift of XRF 050416a by interpolating over the photometric spectrum and stretching the arrival time of the photons by a factor of $(1+z)$.

Shown in Figure 2 are the synthesized light-curves for SN 1998bw at $z = 0.6528$, summed together with the afterglow model. The *HST* data are roughly comparable with the flux normalization and evolution of the summed model. Therefore, the temporal and spectral properties of the *HST* data suggest that XRF 050416a was associated with a supernova similar to SN 1998bw. However, we caution that the temporal coincidence of the SN peak with the radio flare makes it difficult to estimate the relative contributions of the SN and afterglow.

5. Host Galaxy Properties

We now turn to the properties of the GRB host galaxy. We measure the brightness of the host galaxy in the final *HST* epoch to be $F775W = 23.1 \pm 0.1$ mag ($I = 22.7 \pm 0.1$ mag). These values are not corrected for extinction. At $z = 0.6528$ the rest-frame B -band is traced by the observed $F775W$ band, leading to an absolute magnitude, $M_B \approx -20.3 \pm 0.1$ mag, or a luminosity $L_B \approx 0.5L^*$. This host luminosity is similar to that inferred for XRF 030528 (Rau, Salvato & Greiner 2005) and the hosts of typical GRBs (Le Floc’h *et al.* 2003). At

$z = 0.6528$, the measured offset of the optical transient (§2.1) relative to the center of the host galaxy corresponds to 140 ± 140 pc. This offset is a factor of ~ 10 smaller than the median value for long-duration GRBs (Bloom, Kulkarni & Djorgovski 2002).

As shown in Figure 5, the host exhibits several emission lines typical of star-forming galaxies. We estimate the star formation rate in the host galaxy from the observed fluxes of the various emission lines. Using the flux of the [O II] $\lambda 3727$ line, $F_{[\text{O II}]} \approx 9.6 \times 10^{-17}$ erg cm $^{-2}$ s $^{-1}$ (Table 3), and the conversion of Kennicutt (1998), $\text{SFR} = (1.4 \pm 0.4) \times 10^{-41} L_{[\text{O II}]}$ M $_{\odot}$ yr $^{-1}$, we find a star formation rate of about 2.5 ± 0.7 M $_{\odot}$ yr $^{-1}$. From the H β line flux, $F_{\text{H}\beta} \approx 3.7 \times 10^{-17}$ erg cm $^{-2}$ s $^{-1}$, and assuming the Case-B recombination ratio of $F_{\text{H}\alpha}/F_{\text{H}\beta} = 2.87$ and the conversion of Kennicutt (1998), we infer a star formation rate of $\text{SFR} = 7.9 \times 10^{-42} L_{\text{H}\alpha} \approx 1.5 \pm 0.2$ M $_{\odot}$ yr $^{-1}$. Thus, we conclude that the star formation rate (not corrected for extinction) is roughly 2 M $_{\odot}$ yr $^{-1}$. We note that the observed ratio of H γ /H β = 0.3 ± 0.1 compared to the theoretical value of about 0.47 (Osterbrock 1989) suggests a significant extinction correction (factor of ~ 10) following the method of (Calzetti, Kinney & Storchi-Bergmann 1994). We conclude that the star formation rate for the host of XRF 050416a is similar to those inferred for long-duration GRBs (Christensen, Hjorth & Gorosabel 2004; Berger *et al.* 2003a) and at least an order of magnitude larger than that inferred for XRF 020903 (Soderberg *et al.* 2004a; Bersier *et al.* 2006).

The combination of the inferred star formation rate and host luminosity provides a measure of the specific star formation rate. We find a value of $4 \text{ M}_{\odot} \text{ yr}^{-1} L^*/L$ (uncorrected for extinction), which is about a factor of two lower than the mean specific star formation rate for the hosts of long-duration GRBs (Christensen, Hjorth & Gorosabel 2004).

Next, we use the relative strengths of the oxygen and hydrogen emission lines to infer the ionization state and oxygen abundance. The relevant indicators are $R_{23} \equiv \log (F_{[\text{O II}]} + F_{[\text{O III}]} / F_{\text{H}\beta}) \approx 0.74$ and $O_{32} \equiv \log (F_{[\text{O III}]} / F_{[\text{O II}]}) \approx 0.055$. Using the calibrations of McGaugh (1991) and Kobulnicky & Kewley (2004) we find that for the upper branch the metallicity is $12 + \log (\text{O}/\text{H}) \approx 8.6$ while for the lower branch it is about 7.9; the two branches are due to the double-valued nature of R_{23} in terms of metallicity. Thus, the host metallicity of XRF 050416a is $0.2 - 0.8 Z_{\odot}$, larger than that inferred for XRF 060218 (Modjaz *et al.* 2006; Wiersema *et al.* 2007) and comparable to that for XRF 030528 (Rau, Salvato & Greiner 2005). Moreover, this range is somewhat higher than the typical metallicities for GRB hosts, some of which have metallicities that are $\sim 1/10$ solar (e.g., Prochaska *et al.* 2006).

6. Is XRF 050416a a Short Burst?

With a prompt emission duration of just $T_{90} \approx 2.4$ sec (Sakamoto *et al.* 2006), it is interesting to consider XRF 050416a as a member of the short-hard class of gamma-ray bursts, popularly believed to result from the coalescence of neutron stars or black holes (e.g., Eichler *et al.* 1989). Based on the bi-modal BATSE duration distribution, bursts with $T_{90} \gtrsim 2$ seconds are assumed to belong to the long-duration class (Kouveliotou *et al.* 1993), although a decomposition of the overlapping distributions suggests that a small fraction of short-hard bursts (SHBs) have durations longer than this cut-off. The distinction between short-hard and long-duration bursts is further complicated by the detection of soft X-ray tails lasting several seconds following SHBs 050709 and 050724 (Villasenor *et al.* 2005; Barthelmy *et al.* 2005). This suggests that SHBs are not necessarily characterized by a pure hard emission spectrum (Sakamoto 2006). Related to this issue is the use of spectral lags to distinguish between long and short bursts. As discussed by Norris & Bonnell (2006), long-duration bursts typically have longer lags that correlate with isotropic equivalent prompt gamma-ray luminosity (Norris, Marani & Bonnell 2000). On the other hand, SHBs have negligible (or even negative) spectral lags.

In the case of XRF 050416a, the prompt duration places it between the long- and short-duration classes. The low value of E_p may suggest that it belongs to the long-duration class, however, it is becoming clear that hardness cannot be used to reliably distinguish between the two classes (see Nakar 2007). Similarly, while the negative lag inferred for XRF 050416a may suggest a SHB classification, examples do exist of long-duration BATSE bursts with negative lags (Norris & Bonnell 2006).

Afterglow modeling may provide additional clues. The range of values we infer for the XRF 050416a beaming-corrected energies are overall consistent with those of long-duration GRBs including the subclass of sub-energetic bursts. At the same time, they are roughly consistent with the values inferred for SHBs (Fox *et al.* 2005; Berger *et al.* 2005; Soderberg *et al.* 2006a). However, the low circumburst density, $n \sim 10^{-3}$, is a factor of 10 to 10^4 smaller than the typical values inferred for long-duration GRBs (e.g. Panaitescu & Kumar 2002; Yost *et al.* 2003; Chevalier, Li & Fransson 2004) but comparable to those of SHBs (Fox *et al.* 2005; Berger *et al.* 2005; Bloom *et al.* 2006b; Burrows *et al.* 2006; Panaitescu 2006; Soderberg *et al.* 2006a). The observed radio flare and X-ray flattening at 20 days are atypical for both long- and short- duration GRBs and therefore cannot be used to classify this event. However, it is interesting to note that large variations in the circumburst density are more naturally explained in the context of a massive stellar progenitor with interacting stellar winds (Garcia-Segura, Langer & Mac Low 1996; Ramirez-Ruiz *et al.* 2001).

Next is a discussion of the XRF 050416a host galaxy since long- and short- bursts

may also be distinguished by their environments (see Berger 2006 for a recent review). As discussed in §5, the host is a star-forming galaxy with an inferred SFR and metallicity comparable to those of long-duration GRBs. Moreover, XRF 050416a is located near the center and brightest part of its host galaxy. It is thus consistent with the locations of long-duration bursts with respect to their hosts (Bloom, Kulkarni & Djorgovski 2002; Fruchter *et al.* 2006). For comparison, SHBs are typically localized to low SFR hosts with significant old stellar populations at radial offsets up to a factor of 10 larger than those of typical long-duration bursts (Berger *et al.* 2005; Fox *et al.* 2005; Bloom *et al.* 2006a; Soderberg *et al.* 2006a).

Finally, the discovery of an associated Type Ic supernova is perhaps one of the best methods to distinguish between long- and short-bursts. The discovery of several long-duration bursts and X-ray flashes at $z \lesssim 0.3$ in the last few years has firmly established that GRBs and XRFs are accompanied by supernovae of Type Ic (Galama *et al.* 1998; Stanek *et al.* 2003; Malesani *et al.* 2004; Soderberg *et al.* 2005b; Pian *et al.* 2006). A study of the luminosity distribution for GRB/XRF-SNe reveals a significant dispersion, implying a spread of (at least) an order of magnitude in peak optical luminosity (Zeh, Klose & Hartmann 2004; Soderberg *et al.* 2006d). At the same time, deep imaging of SHBs has constrained any associated SN emission to be up to ~ 100 times less luminous as SN 1998bw (Fox *et al.* 2005; Soderberg *et al.* 2006a). In the case of XRF 050416a, the temporal and spectral properties of the *HST* data suggest that XRF 050416a was accompanied by a supernova with a peak luminosity roughly similar to SN 1998bw. We stress, however, that the temporal coincidence of the radio flare with the SN peak complicates any study of the SN properties.

Based on the associated SN and large-scale environmental properties, we conclude that XRF 050416a is a member of the long-duration class of gamma-ray bursts. This event highlights the difficulty in classifying bursts based on their prompt emission properties alone.

7. Discussion and Conclusions

We present extensive broad-band NIR/optical and radio data for the afterglow of the X-ray flash 050416a and show that it is localized to a star-forming ($\gtrsim 2 M_{\odot} \text{ yr}^{-1}$) host galaxy at $z = 0.6528$. Along with XRFs 020903, 030723 and 060218, this burst is one of the best studied X-ray flashes to date. Moreover, XRF 050416a is only the third XRF with a spectroscopic redshift for which a broadband afterglow study has been performed and the physical parameters have been constrained. The isotropic-equivalent prompt and kinetic energy releases are $E_{\gamma,\text{iso}} \approx 1.2 \times 10^{51}$ (Sakamoto *et al.* 2006) and $E_{K,\text{iso}} \approx 10^{52}$ erg respectively. These values are a factor of 10^2 times larger than those of XRF 020903 (Sakamoto *et al.*

2004; Soderberg *et al.* 2004a) and up to 10^4 times larger than those inferred for sub-energetic GRBs 980425, 031203 and XRF 060218 (Pian *et al.* 2000; Kulkarni *et al.* 1998; Sazonov, Lutovinov & Sunyaev 2004; Soderberg *et al.* 2004b; Campana *et al.* 2006; Soderberg *et al.* 2006c).

Adopting the results of our standard synchrotron model and $t_j \gtrsim 20$ days we constrain the collimation of the ejecta to $\theta_j \gtrsim (6.9 - 9.7)^\circ$, slightly larger than the median value of 5° for typical GRBs (Bloom, Frail & Kulkarni 2003; Ghirlanda, Ghisellini & Lazzati 2004 and references therein). This indicates that the beaming-corrected energy release is $E_K \approx (9.8 \times 10^{49} - 1.4 \times 10^{52})$ erg and $E_\gamma \approx (8.6 \times 10^{48} - 1.2 \times 10^{51})$ erg, implying a total relativistic energy yield of $E_{\text{tot}} \approx (1.1 \times 10^{50} - 1.5 \times 10^{52})$ erg, which straddles the median value for cosmological GRBs, $E_{\text{tot}} \approx 2 \times 10^{51}$ erg (Berger *et al.* 2003b). However, the efficiency in converting the energy in the ejecta into γ -rays is $\eta_\gamma \equiv E_\gamma / (E_K + E_\gamma) \approx (0.04 - 0.08)$, significantly lower than the typical values for GRBs (Panaitescu & Kumar 2002; Yost *et al.* 2003) and comparable to that inferred for XRF 020903 ($\eta_\gamma \approx 0.03$, Soderberg *et al.* 2004a). This strengthens the idea that XRFs and GRBs are distinguished by their ability to couple significant energy to highly-relativistic material (Soderberg *et al.* 2004a; Zhang, Woosley & Heger 2004).

In addition to the burst energetics, several key results emerge from our broadband analysis of XRF 050416a. First is the detection of a bright, late-time radio flare accompanied by an observed flattening in the X-ray bands which we attribute to a large circumburst density enhancement or episode of energy injection (refreshing shell or off-axis ejecta) at $t \sim 20$ days. In the context of a density jump, it is interesting to note that the inferred pre-jump circumburst density is several orders of magnitude lower than the values typically inferred for long-duration GRBs (see Soderberg *et al.* 2006e for a recent compilation). Moreover, radio observations of local (optically-selected) core-collapse SNe show similar flux modulations (factor of 2 to 3) attributed to abrupt variations in the circumstellar medium (Weiler *et al.* 1991; Ryder *et al.* 2004; Soderberg *et al.* 2006b).

Next, the temporal and spectral evolution of the optical afterglow suggests the contribution from a supernova component with peak luminosity and light-curve shape comparable to SN 1998bw. Given the temporal coincidence of the radio flare and X-ray flattening with the SN peak, XRF 050416a highlights the need for full spectral coverage in late-time GRB/XRF-SN searches. This is illustrated by the optical rebrightening observed for XRF 030723 at ~ 15 days and interpreted as a thermal SN component (Fynbo *et al.* 2004) while the X-ray and radio data show similar rebrightenings at late-time suggesting a CSM density jump or energy injection (Butler *et al.* 2005, Soderberg *et al.* in prep).

With the addition of XRF 050416a, there are three XRFs (020903, 060218 and now 050416a) with spectroscopic redshifts observed in association with supernovae with peak

luminosities varying by up to a factor of 10 compared to SN 1998bw (Soderberg *et al.* 2005b; Bersier *et al.* 2006; Pian *et al.* 2006). However, deep *HST* observations of XRF 040701 at $z = 0.21$ revealed that any associated SN was at least a factor of 10 (and likely $\gtrsim 100$) times fainter than SN 1998bw (Soderberg *et al.* 2005b). It is therefore clear that most XRFs produce Nickel-rich supernova explosions, but that there is a significant dispersion in the peak luminosities of XRF-associated SNe. A similar result is found for GRB-associated SNe (Zeh, Klose & Hartmann 2004; Soderberg *et al.* 2006d) and further strengthens the idea that GRBs and XRFs are intimately related.

Finally, we address XRF 050416a within the framework of off-axis and dirty fireball models for XRFs. As discussed in §3.3, the evolution of the early afterglow indicates that the ejecta are being viewed along the collimation axis. Similarly, on-axis viewing angles are inferred from afterglow studies of XRFs 020903 (Soderberg *et al.* 2004a), 050215B (Levan *et al.* 2006), 050406 (Romano *et al.* 2006), and 060218 (Campana *et al.* 2006; Soderberg *et al.* 2006c). Unification models in which X-ray flashes are understood as typical GRBs viewed away from the burst collimation axis (e.g. Granot *et al.* 2002; Yamazaki, Yonetoku & Nakamura 2003) are therefore inconsistent with the observations for these five events. To date, such models are only consistent with the observations of one X-ray flash: XRF 030723 (Fynbo *et al.* 2004; Butler *et al.* 2005; Granot, Ramirez-Ruiz & Perna 2005).

While the lack of high energy photons prevents a direct constraint on the Lorentz factor of the burst, our detailed afterglow modeling allows us to place a lower limit on the initial bulk Lorentz factor, $\Gamma_0 \gtrsim 110$, by extrapolating $\Gamma \propto t^{-3/8}$ back to the first XRT observation at ~ 100 sec. This value is somewhat lower than the typical values inferred for cosmological GRBs (Panaiteescu & Kumar 2002) and consistent with the upper range of values predicted for a dirty fireball. Given that some dirty fireball models predict XRFs to have wider jets than typical GRBs (e.g., Zhang, Woosley & MacFadyen 2003), it is interesting to note that there are no X-ray flashes for which an achromatic jet break has been observed. This may suggest that their ejecta are not strongly collimated. In fact, detailed radio observations of XRFs 020903 and 060218 spanning $\gtrsim 100$ days imply a quasi-spherical ejecta geometry in both cases (Soderberg *et al.* 2004a, 2006c).

While the basic properties of X-ray flashes (redshifts, hosts, isotropic γ -ray energies) are now available for several events, it is clear that broadband afterglow observations are required for a complete understanding of the burst properties. With the addition of XRF 050416a to the existing sample of only a few well-studied events, we continue to work towards a systematic comparison of the global characteristics of XRFs and GRBs. While limited, the current sample suggests that the two classes share similar ejecta energies, associated SNe, and viewing angles. Further studies of XRF afterglows will be used to confirm whether the

two classes differ in their collimation angles.

The authors thank Re'em Sari and Don Lamb for helpful discussions. As always, the authors thank Jochen Greiner for maintaining his GRB page. A.M.S. and S.B.C. are supported by the NASA Graduate Student Research Program. E.B. and A.G. acknowledge support by NASA through Hubble Fellowship grants awarded by STScI, which is operated by AURA, Inc., for NASA. GRB research at Caltech is supported through NASA.

REFERENCES

- Alard, C. 2000, *A&AS*, 144, 363.
- Amati, L. 2006, *MNRAS*, 2006, 372, 233.
- Band, D. *et al.* 1993, *ApJ*, 413, 281.
- Barthelmy, S. D. *et al.* 2005, *Nature*, 438, 994.
- Berger, E. 2006, in *AIP Conf. Proc. 838: Gamma-Ray Bursts in the Swift Era*, ed. S. S. Holt, N. Gehrels, and J. A. Nousek, 33.
- Berger, E., Cowie, L. L., Kulkarni, S. R., Frail, D. A., Aussel, H., and Barger, A. J. 2003a, *ApJ*, 588, 99.
- Berger, E., Kulkarni, S. R., and Frail, D. A. 2003, *ApJ*, 590, 379.
- Berger, E., Kulkarni, S. R., and Frail, D. A. 2004, *ApJ*, 612, 966.
- Berger, E. *et al.* 2003b, *Nature*, 426, 154.
- Berger, E. *et al.* 2005, *Nature*, 438, 988.
- Bersier, D. *et al.* 2006, *ApJ*, 643, 284.
- Björnsson, G., Gudmundsson, E. H., and Jóhannesson, G. 2004, *ApJ*, 615, L77.
- Bloom, J. S., Fox, D., van Dokkum, P. G., Kulkarni, S. R., Berger, E., Djorgovski, S. G., and Frail, D. A. 2003, *ApJ*, 599, 957.
- Bloom, J. S., Frail, D. A., and Kulkarni, S. R. 2003, *ApJ*, 594, 674.
- Bloom, J. S., Kulkarni, S. R., and Djorgovski, S. G. 2002, *AJ*, 123, 1111.
- Bloom, J. S. *et al.* 2006a, *astro-ph/0607223*

- Bloom, J. S. *et al.* 2006b, ApJ, 638, 354.
- Burrows, D. N. *et al.* 2006, ApJ, 653, 468.
- Butler, N. R. *et al.* 2005, ApJ, 621, 884.
- Calzetti, D., Kinney, A. L., and Storchi-Bergmann, T. 1994, ApJ, 429, 582.
- Campana, S. *et al.* 2006, Nature, 442, 1008.
- Chevalier, R. A., Li, Z.-Y., and Fransson, C. 2004, ApJ, 606, 369.
- Christensen, L., Hjorth, J., and Gorosabel, J. 2004, A&A, 425, 913.
- Dermer, C. D., Chiang, J., and Böttcher, M. 1999, ApJ, 513, 656.
- Eichler, D., Livio, M., Piran, T., and Schramm, D. N. 1989, Nature, 340, 126.
- Fox, D. B. *et al.* 2005, Nature, 437, 845.
- Frail, D. A., Kulkarni, S. R., Berger, E., and Wieringa, M. H. 2003, AJ, 125, 2299.
- Frail, D. A. *et al.* 2001, ApJ, 562, L55.
- Frail, D. A., Soderberg, A. M., Kulkarni, S. R., Berger, E., Yost, S., Fox, D. W., and Harrison, F. A. 2005, ApJ, 619, 994.
- Frail, D. A., Waxman, E., and Kulkarni, S. R. 2000, ApJ, 537, 191.
- Freedman, D. L. and Waxman, E. 2001, ApJ, 547, 922.
- Fruchter, A. S. and Hook, R. N. 2002, PASP, 114, 144.
- Fruchter, A. S. *et al.* 2006, Nature, 441, 463.
- Fynbo, J. P. U. *et al.* 2004, ApJ, 609, 962.
- Galama, T. J. *et al.* 1998, Nature, 395, 670.
- Galama, T. J. and Wijers, R. A. M. J. 2001, ApJ, 549, L209.
- Garcia-Segura, G., Langer, N., and Mac Low, M.-M. 1996, A&A, 316, 133.
- Ghirlanda, G., Ghisellini, G., and Lazzati, D. 2004, ApJ, 616, 331.
- Granot, J., Nakar, E., and Piran, T. 2003, Nature, 426, 138.

- Granot, J., Panaitescu, A., Kumar, P., and Woosley, S. E. 2002, *ApJ*, 570, L61.
- Granot, J. and Sari, R. 2002, *ApJ*, 568, 820.
- Granot, J. and Ramirez-Ruiz, E. and Perna, R. 2005, *ApJ*, 630, 1003.
- Heise, J., in’t Zand, J., Kippen, R. M., and Woods, P. M. 2001, in *Gamma-ray Bursts in the Afterglow Era*, ed. E. Costa, F. Frontera, and J. Hjorth, 16.
- Henden, A. 2005, *GRB Coordinates Network*, 3454, 1.
- Holland, S. T. *et al.* 2006, *ArXiv Astrophysics e-prints*. astro-ph/0604316.
- Jakobsson, P. *et al.* 2004, *A&A*, 427, 785.
- Kennicutt, R. C. 1998, *ARA&A*, 36, 189.
- Kobulnicky, H. A. and Kewley, L. J. 2004, *ApJ*, 617, 240.
- Kouveliotou, C., Meegan, C. A., Fishman, G. J., Bhat, N. P., Briggs, M. S., Koshut, T. M., Paciesas, W. S., and Pendleton, G. N. 1993, *ApJ*, 413, L101.
- Kouveliotou, C. *et al.* 2004, *ApJ*, 608, 872.
- Kulkarni, S. R. *et al.* 1999, *ApJ*, 522, L97.
- Kulkarni, S. R. *et al.* 1998, *Nature*, 395, 663.
- Le Floch, E. *et al.* 2003, *A&A*, 400, 499.
- Levan, A. J. *et al.* 2006, *ApJ*, 648, 1132.
- Li, W., Chornock, R., Jha, S., and Filippenko, A. V. 2005, *GRB Coordinates Network*, 3270, 1.
- Li, Z. and Song, L. M. 2004, *ApJ*, 614, L17.
- Lithwick, Y. and Sari, R. 2001, *ApJ*, 555, 540.
- Livio, M. and Waxman, E. 2000, *ApJ*, 538, 187.
- Malesani, D. *et al.* 2004, *ApJ*, 609, L5.
- Mangano, V. *et al.* 2006, *ArXiv Astrophysics e-prints*. astro-ph/0603738.
- McGaugh, S. S. 1991, *ApJ*, 380, 140.

- McKenzie, E. H. and Schaefer, B. E. 1999, PASP, 111, 964.
- Mirabal, N., Halpern, J. P., An, D., Thorstensen, J. R., and Terndrup, D. M. 2006, ApJ, 643, L99.
- Modjaz, M. *et al.* 2006, ApJ, 645, L21.
- Nakar, E. and Granot, J. 2006, ArXiv Astrophysics e-prints. astro-ph/0606011.
- Nakar, E. 2007, Physics Reports (in press). astro-ph/0701748.
- Norris, J. P. and Bonnell, J. T. 2006, ApJ, 643, 266.
- Norris, J. P., Marani, G. F., and Bonnell, J. T. 2000, ApJ, 534, 248.
- Nousek, J. A. *et al.* 2006, ApJ, 642, 389.
- O’Brien, P. T. *et al.* 2006, ApJ, 647, 1213.
- Osterbrock, D. E. 1989, Astrophysics of gaseous nebulae and active galactic nuclei, (University of Minnesota, et al. Mill Valley, CA: Research supported by the University of California, John Simon Guggenheim Memorial Foundation, University Science Books).
- Paczynski, B. 2001, Acta Astronomica, 51, 1.
- Panaiteanu, A. 2006, MNRAS, 367, L42.
- Panaiteanu, A. and Kumar, P. 2002, ApJ, 571, 779.
- Patat, F. *et al.* 2001, ApJ, 555, 900.
- Pei, Y. C. 1992, ApJ, 395, 130.
- Pian, E. *et al.* 2000, ApJ, 536, 778.
- Pian, E. *et al.* 2006, Nature, 442, 1011.
- Piran, T. 1999, Phys. Rep., 314, 575.
- Predehl, P. and Schmitt, J. H. M. M. 1995, A&A, 293, 889.
- Price, P. A., Minezaki, T., Cowie, L., and Yoshii, Y. 2005, GRB Coordinates Network, 3312, 1.
- Prochaska, J. X. *et al.* 2006, ApJ, 642, 989.

- Qiu, Y., Lu, C. L., Lou, Y. Q., Huang, K. Y., and Urata, Y. 2005, GRB Coordinates Network, 3286, 1.
- Ramirez-Ruiz, E., Dray, L. M., Madau, P., and Tout, C. A. 2001, MNRAS, 327, 829.
- Ramirez-Ruiz, E., García-Segura, G., Salmonson, J. D., and Pérez-Rendón, B. 2005, ApJ, 631, 435.
- Rau, A., Salvato, M., and Greiner, J. 2005, A&A, 444, 425.
- Rees, M. J. and Meszaros, P. 1998, ApJ, 496, L1+.
- Romano, P. *et al.* 2006, A&A, 450, 59.
- Ryder, S. D., Sadler, E. M., Subrahmanyam, R., Weiler, K. W., Panagia, N., and Stockdale, C. 2004, MNRAS, 349, 1093.
- Sakamoto, T. 2006, in AIP Conf. Proc. 838: Gamma-Ray Bursts in the Swift Era, ed. S. S. Holt, N. Gehrels, and J. A. Nousek, 43.
- Sakamoto, T. *et al.* 2006, ApJ, 636, L73.
- Sakamoto, T. *et al.* 2004, ApJ, 602, 875.
- Sakamoto, T. *et al.* 2005, ApJ, 629, 311.
- Sari, R. and Piran, T. 1999, ApJ, 517, L109.
- Sari, R., Piran, T., and Halpern, J. P. 1999, ApJ, 519, L17.
- Sari, R., Piran, T., and Narayan, R. 1998, ApJ, 497, L17+.
- Sazonov, S. Y., Lutovinov, A. A., and Sunyaev, R. A. 2004, Nature, 430, 646.
- Schady, P. *et al.* 2006, ApJ, 643, 276.
- Schlegel, D. J., Finkbeiner, D. P., and Davis, M. 1998, ApJ, 500, 525.
- Sirianni, M. *et al.* 2005, PASP, 117, 1049.
- Smith, J. A. *et al.* 2002, AJ, 123, 2121.
- Soderberg, A. M. 2006, in AIP Conf. Proc. 838: Gamma-Ray Bursts in the Swift Era, ed. S. S. Holt, N. Gehrels, and J. A. Nousek, 380.
- Soderberg, A. M. *et al.* 2006a, ApJ, 650, 261.

- Soderberg, A. M., Chevalier, R. A., Kulkarni, S. R., and Frail, D. A. 2006b, *ApJ*, 651, 1005.
- Soderberg, A. M. *et al.* 2004a, *ApJ*, 606, 994.
- Soderberg, A. M. *et al.* 2004b, *Nature*, 430, 648.
- Soderberg, A. M. *et al.* 2005b, *ApJ*, 627, 877.
- Soderberg, A. M. *et al.* 2006c, *Nature*, 442, 1014.
- Soderberg, A. M. *et al.* 2006d, *ApJ*, 636, 391.
- Soderberg, A. M., Nakar, E., Berger, E., and Kulkarni, S. R. 2006e, *ApJ*, 638, 930.
- Stanek, K. Z. *et al.* 2003, *ApJ*, 591, L17.
- Tiengo, A., Mereghetti, S., Ghisellini, G., Rossi, E., Ghirlanda, G., and Schartel, N. 2003, *A&A*, 409, 983.
- Torii, K. 2005, *GRB Coordinates Network*, 3272, 1.
- Villasenor, J. S. *et al.* 2005, *Nature*, 437, 855.
- Waxman, E. 2004, *ApJ*, 602, 886.
- Weiler, K. W., van Dyk, S. D., Discenna, J. L., Panagia, N., and Sramek, R. A. 1991, *ApJ*, 380, 161.
- Wiersema, K. *et al.* 2007, *A&A*, in press (astro-ph/0701034).
- Wijers, R. A. M. J. 2001, in *Gamma-ray Bursts in the Afterglow Era*, ed. E. Costa, F. Frontera, and J. Hjorth, 306.
- Yamazaki, R., Yonetoku, D., and Nakamura, T. 2003, *ApJ*, 594, L79.
- Yanagisawa, K., Toda, H., and Kawai, N. 2005, *GRB Coordinates Network*, 3287, 1.
- Yost, S. A., Harrison, F. A., Sari, R., and Frail, D. A. 2003, *ApJ*, 597, 459.
- Zeh, A., Klose, S., and Hartmann, D. H. 2004, *ApJ*, 609, 952.
- Zel’dovich, Y. B. and Raizer, Y. P. 2002, *Physics of Shock Waves and High Temperature Hydrodynamic Phenomena*, (Mineola, NY: Dover).
- Zhang, W., Woosley, S. E., and Heger, A. 2004, *ApJ*, 608, 365.

Zhang, W., Woosley, S. E., and MacFadyen, A. I. 2003, ApJ, 586, 356.

Table 1. Ground-based Optical and NIR Observations of XRF 050416a

Date Obs (UT)	Δt^a (days)	Telescope	Filter	magnitude ^b
2005 April 16.4641	0.0025	Palomar 60-inch	<i>I</i>	18.82 ± 0.11
2005 April 16.4659	0.0043	Palomar 60-inch	<i>I</i>	18.86 ± 0.11
2005 April 16.4677	0.0061	Palomar 60-inch	<i>I</i>	19.16 ± 0.13
2005 April 16.4696	0.0080	Palomar 60-inch	<i>I</i>	19.35 ± 0.23
2005 April 16.4692	0.0076	Palomar 200-inch	<i>K_s</i>	16.37 ± 0.21
2005 April 16.4714	0.0098	Palomar 60-inch	<i>I</i>	19.01 ± 0.12
2005 April 16.4731	0.0116	Palomar 60-inch	<i>I</i>	19.53 ± 0.18
2005 April 16.4772	0.0156	Palomar 60-inch	<i>z'</i>	19.16 ± 0.27^c
2005 April 16.4763	0.0147	Palomar 200-inch	<i>K_s</i>	16.64 ± 0.20
2005 April 16.4833	0.0217	Palomar 60-inch	<i>I</i>	19.09 ± 0.20
2005 April 16.4828	0.0212	Palomar 200-inch	<i>K_s</i>	16.79 ± 0.20
2005 April 16.4880	0.0264	Palomar 60-inch	<i>I</i>	19.34 ± 0.23
2005 April 16.4895	0.0279	Palomar 200-inch	<i>K_s</i>	16.89 ± 0.24
2005 April 16.4960	0.0344	Palomar 200-inch	<i>K_s</i>	16.99 ± 0.25
2005 April 16.5026	0.0410	Palomar 200-inch	<i>K_s</i>	16.98 ± 0.29
2005 April 16.5089	0.0473	ANU 2.3-meter	<i>R</i>	20.92 ± 0.05
2005 April 16.5502	0.0886	ANU 2.3-meter	<i>R</i>	21.18 ± 0.12
2005 April 16.5543	0.0927	ANU 2.3-meter	<i>R</i>	21.40 ± 0.15
2005 April 16.5585	0.0969	ANU 2.3-meter	<i>R</i>	21.16 ± 0.08
2005 April 16.5626	0.1010	ANU 2.3-meter	<i>R</i>	21.43 ± 0.11
2005 April 16.5669	0.1053	ANU 2.3-meter	<i>R</i>	21.44 ± 0.26
2005 April 16.5702	0.1086	ANU 2.3-meter	<i>R</i>	21.39 ± 0.16
2005 April 16.5744	0.1128	ANU 2.3-meter	<i>R</i>	21.37 ± 0.16
2005 April 16.5779	0.1163	ANU 2.3-meter	<i>R</i>	21.18 ± 0.17
2005 April 16.5816	0.1200	ANU 2.3-meter	<i>R</i>	21.35 ± 0.16
2005 April 16.5850	0.1234	ANU 2.3-meter	<i>R</i>	21.36 ± 0.10
2005 April 16.5893	0.1277	ANU 2.3-meter	<i>R</i>	21.44 ± 0.27
2005 April 16.5924	0.1308	ANU 2.3-meter	<i>R</i>	21.46 ± 0.22
2005 April 16.5975	0.1359	ANU 2.3-meter	<i>R</i>	21.63 ± 0.34
2005 April 16.6092	0.1476	ANU 2.3-meter	<i>R</i>	21.71 ± 0.30

Table 1—Continued

Date Obs (UT)	Δt^a (days)	Telescope	Filter	magnitude ^b
2005 April 16.6187	0.1571	ANU 2.3-meter	<i>R</i>	21.87 ± 0.46
2005 April 17.3895	0.9279	Palomar 60-inch	<i>R</i>	22.62 ± 0.30
2005 April 17.4006	0.9390	Palomar 60-inch	<i>z'</i>	$< 20.38^c$
2005 April 17.4087	0.9467	Palomar 60-inch	<i>I</i>	22.05 ± 0.30
2005 April 18.2884	1.8264	Palomar 60-inch	<i>R</i>	< 23.46
2005 April 18.4031	1.9411	Palomar 60-inch	<i>I</i>	< 22.83
2005 April 23.3511	6.8896	Palomar 200-inch	<i>K_s</i>	< 19.70

^aDays since explosion have been calculated for the mid-point of each exposure.

^bMagnitudes have not been corrected for extinction. Limits are given as 3σ .

^cAB system.

Table 2. *HST/ACS* Observations of XRF 050416a

Date Obs (UT)	Δt (days)	Exp. Time (sec)	Filter	<i>HST</i> mag ^a (AB)	Johnson mag ^b (Vega)
2005 May 23.38	36.92	3282	F775W	24.35 ± 0.02	23.82 ± 0.02
2005 May 23.46	37.00	3430	F850LP	23.83 ± 0.03	...
2005 June 16.17	60.71	3986	F775W	25.88 ± 0.10	25.36 ± 0.10
2005 July 11.15	85.69	4224	F775W	26.44 ± 0.22	25.92 ± 0.22
2005 November 21.06	218.60	4224	F850LP	...	...
2005 November 21.19	218.73	4224	F775W	...	...

^aAB system magnitudes in the *HST* filters given in column 4. Photometry was done on residual images (see §2.2). We have assumed the source flux to be negligible in the final (template) epoch. Magnitudes have not been corrected for extinction.

^bWe convert the F775W magnitudes in column 5 to Johnson *I*-band (Vega system) as described in §2.2. Magnitudes have not been corrected for extinction.

Table 3. Spectroscopic lines for XRF 050416a

Line	λ (rest) (Å)	λ (observed) ^a (Å)	Redshift	Flux ^b ($\times 10^{-17}$ erg cm ⁻² s ⁻¹)
[OII]	3728.38	6162.08	0.6528	9.6 ± 0.4
H γ	4341.72	7174.64	0.6525	1.2 ± 0.2
H β	4862.72	8037.14	0.6528	3.7 ± 0.5
[OIII]	4960.30	8198.79	0.6529	2.5 ± 0.3
[OIII]	5008.24	8277.57	0.6528	8.4 ± 0.4

^aObserved wavelengths have been corrected to vacuum.

^bFlux values have not been corrected for Galactic extinction

Table 4. Radio Observations of XRF 050416a

Date Obs (UT)	Δt (days)	$F_{\nu, 1.43 \text{ GHz}}^{\text{a}}$ (μJy)	$F_{\nu, 4.86 \text{ GHz}}$ (μJy)	$F_{\nu, 8.46 \text{ GHz}}$ (μJy)
2005 April 16.49	0.026	...	...	20 ± 51
2005 April 22.03	5.57	...	200 ± 46	101 ± 34
2005 April 28.29	11.83	...	188 ± 42	132 ± 30
2005 May 1.35	14.89	...	201 ± 43	...
2005 May 31.10	44.64	...	...	431 ± 46
2005 June 3.99	48.53	...	585 ± 48	...
2005 June 8.03	52.57	420 ± 190	562 ± 51	398 ± 33
2005 June 16.97	61.51	...	505 ± 49	330 ± 35
2005 June 20.16	64.70	...	...	286 ± 33
2005 July 1.92	76.46	843 ± 214	321 ± 48	189 ± 36
2005 July 30.02	104.56	0 ± 192	68 ± 36	93 ± 41
2005 August 14.97	120.51	...	...	72 ± 42
2005 August 21.89	127.42	...	...	41 ± 47
2005 September 19.90	156.44	...	0 ± 54	...
2005 October 15.73	182.27	...	33 ± 47	89 ± 38

^aAll errors are given as 1σ (rms).




Fig. 1.— Discovery image of the afterglow of XRF 050416a. We began observing the field of XRF 050416a with the Palomar 60-inch telescope in the I -band about 2.5 minutes after the burst. We discovered a fading source within the 3.0 arcmin (radius) *Swift*/BAT error circle. Subsequent localizations of the XRT (circle) and UVOT afterglow positions were shown to be coincident with the P60 source. Our late-time *HST* images reveal a host galaxy coincident with the optical afterglow position.




Fig. 2.— Optical ($BVRIz'$) and NIR (K_s) light-curves of the afterglow of XRF 050416a. We supplement our measurements from Table 1 (black symbols) with data from the GCNs (Li *et al.* 2005; Price *et al.* 2005; Qiu *et al.* 2005; Torii 2005; Yanagisawa, Toda & Kawai 2005) and Holland *et al.* (2006) (grey symbols). Detections are shown as circles and upper limits as inverted triangles. R -band data from the ANU 2.3-meter have been binned for clarity. All data have been corrected for host galaxy extinction (see §2.1). Between $t \approx 0.01$ and 1 day the NIR/optical afterglow shows an average decay index of $\overline{\alpha}_{\text{NIR/opt}} \approx -0.75$. As described in §3.2, we find a reasonable fit to the spectral and temporal evolution of the broadband data with a standard afterglow model. Our *HST* measurements are shown as encircled dots; the implied steep spectral index suggests the contribution from an supernova component at $t \sim 40$ days (§4.4). We sum the flux from a SN 1998bw-like supernova at $z = 0.6528$ with our afterglow model to produce the final fits (grey solid lines). For comparison, we also show the afterglow model alone (dashed grey line). We note that this model does not apply to the very early afterglow evolution at $t \lesssim 0.01$ days. The dotted grey lines at early time represent the evolution of the NIR/optical assuming these bands track the X-ray evolution on this timescale.




Fig. 3.— The observed photometric spectrum for XRF050416a is shown at $t \approx 0.014$ (top) and 0.8 days (bottom). The K_s –, z' –, I – and R –band data (black symbols) are from Table 1 while the V , B , U , $UVW1$, $UVM2$ and $UVW2$ data (grey symbols) are from Holland *et al.* (2006). Detections are shown as circles and upper limits as inverted triangles. The observed spectrum is fit by $\beta_{\text{NIR/opt}} \approx -1.3$ and -1.5 in the top and bottom panels, respectively (dotted line). We fit the data with an extinction model (solid line) which includes $E(B - V) = 0.03$ from the Galaxy (Schlegel, Finkbeiner & Davis 1998) and a rest-frame host galaxy extinction of $E(B - V)_{\text{rest}} = 0.28$ assuming an intrinsic spectral index of $\beta_{\text{NIR/opt}} \approx -0.55$ (dashed line).




Fig. 4.— *HST* images of XRF 050416a were obtained with *ACS* in filters F775W and F850LP spanning $t \approx 37$ to 219 days after the explosion. The host galaxy emission dominates that of the afterglow throughout this timescale. Relative astrometry between the P60 and the *HST* images provides an optical afterglow position accurate to 0.70 arcsec (circle, 2σ). Residual images of the afterglow were produced through image subtraction techniques (§2.2). The position of the afterglow is offset by 0.02 ± 0.02 arcsec with respect to the host galaxy center.




Fig. 5.— Optical spectrum of the host galaxy of XRF 050416a as observed with the Low Resolution Imaging Spectrograph on Keck I on 2005 June 6.3 UT. Several bright emission lines are detected, indicating that the host is a star-forming galaxy at redshift $z = 0.6528 \pm 0.0002$. As discussed in §5 we find that the host is actively forming stars with a rate of $\gtrsim 2 M_{\odot} \text{ yr}^{-1}$ and an inferred metallicity of $Z \sim 0.2 - 0.8 Z_{\odot}$.




Fig. 6.— Radio observations of XRF 050416a from the Very Large Array as listed in Table 4. We have summed the late-time limits in each frequency for clarity. The radio evolution is characterized by an unusual flare at $t \sim 40$ days which rapidly fades below our detection limits. This radio flare occurs roughly on the same timescale as the observed X-ray flattening. (Figure 7). We attribute both the radio flare and the X-ray flattening to an episode of late-time energy injection (refreshing shell or off-axis ejecta) or a large circumburst density jump. Our early afterglow model fit is shown (solid grey lines) and extrapolated to the epoch of the radio flare (dashed grey lines). The spectral index between 8.46 and 4.86 GHz is optically thin throughout our radio monitoring, including the late-time flare (lower right panel).




Fig. 7.— X-ray observations of XRF 050416a from the *Swift*/XRT as reported by Mangano *et al.* (2006). The X-ray light-curve is characterized by four phases: (1) a initial steep decay, (2) a flattening between $t \approx 0.005 - 0.01$ days, (3) a subsequent decay, and (4) a second flattening between $t \approx 20 - 40$ days. The final measurement at 70 days implies a rapid steepening following the second flattening. We attribute the late-time flattening to energy injection (refreshing shell or off-axis ejecta) or a large circumburst density jump. Overplotted is our afterglow model fit between 0.014 and 20 days (solid grey line) and extrapolated to late time (dashed grey line). We do not attempt to fit the data prior to 0.01 days (dotted line) due to insufficient broadband data.